

# Economics 765

May 26, 2017

R. Davidson

## Assignment 3

You are asked to do exercises 3.2, 3.7, 4.5, and 4.13 of Volume 2 of Shreve. The essence of these exercises is reproduced below for convenience.

**3.2** Let  $W(t)$ ,  $t \geq 0$ , be a Brownian motion, and let  $\mathcal{F}(t)$ ,  $t \geq 0$ , be a filtration for this Brownian motion. Show that  $W^2(t) - t$  is a martingale.

**3.7** Theorem 3.6.2 provides the so-called *Laplace transform* of the density of the first passage time for Brownian motion (the moment-generating function). Let  $W$  be a Brownian motion. Fix  $m > 0$  and  $\mu \in \mathbb{R}$ . For  $0 \leq t < \infty$ , define

$$\begin{aligned} X(t) &= \mu t + W(t), \\ \tau_m &= \min\{t \geq 0; X(t) = m\}. \end{aligned}$$

As usual, we set  $\tau_m = \infty$  if  $X(t)$  never reaches the level  $m$ . Let  $\sigma$  be a positive number and set

$$Z(t) = \exp \left\{ \sigma X(t) - \left( \sigma \mu + \frac{1}{2} \sigma^2 \right) t \right\}.$$

(i) Show that  $Z(t)$ ,  $t \geq 0$ , is a martingale.

(ii) Use (i) to conclude that

$$\mathbb{E} \left[ \exp \left\{ \sigma X(t \wedge \tau_m) - \left( \sigma \mu + \frac{1}{2} \sigma^2 \right) (t \wedge \tau_m) \right\} \right] = 1, \quad t \geq 0.$$

(iii) Now suppose  $\mu \geq 0$ . Show that, for  $\sigma > 0$ ,

$$\mathbb{E} \left[ \exp \left\{ \sigma m - \left( \sigma \mu + \frac{1}{2} \sigma^2 \right) \tau_m \right\} \mathbf{I}(\tau_m < \infty) \right] = 1.$$

Use this fact to show that  $P\{\tau_m < \infty\} = 1$  and to obtain the Laplace transform

$$\mathbb{E} e^{-\alpha \tau_m} = e^{m\mu - m\sqrt{2\alpha + \mu^2}} \quad \text{for all } \alpha > 0.$$

(iv) Show that, if  $\mu > 0$ , then  $\mathbb{E}\tau_m < \infty$ . Obtain a formula for  $\mathbb{E}\tau_m$ . (Hint: Differentiate the formula in (ii) with respect to  $\alpha$ .)

(v) Now suppose  $\mu < 0$ . Show that, for  $\sigma > -2\mu$ ,

$$\mathbb{E} \left[ \exp \left\{ \sigma m - \left( \sigma \mu + \frac{1}{2} \sigma^2 \right) \tau_m \right\} \mathbf{I}(\tau_m < \infty) \right] = 1.$$

Use this fact to show that  $P\{\tau_m < \infty\} = e^{-2m|\mu|}$  (watch out! there is a misprint here in Shreve, who writes  $e^{-2x|\mu|}$ ), which is strictly less than 1, and to obtain the Laplace transform

$$\mathbb{E} e^{-\alpha \tau_m} = e^{m\mu - m\sqrt{2\alpha + \mu^2}} \quad \text{for all } \alpha > 0.$$

**4.5** Let  $S(t)$  be a positive stochastic process that satisfies the generalised geometric Brownian motion differential equation

$$dS(t) = \alpha(t)S(t) dt + \sigma(t)S(t) dW(t),$$

where  $\alpha(t)$  and  $\sigma(t)$  are processes adapted to the filtration  $\mathcal{F}(t)$ ,  $t \geq 0$ , associated with the Brownian motion  $W(t)$ ,  $t \geq 0$ .

- (i) Make use of the above differential equation and the It-Doebelin formula in order to compute  $d \log S(t)$ . Simplify so that you have a formula for  $d \log S(t)$  that does not involve  $S(t)$ .
- (ii) Integrate the formula you obtained in (i), and then exponentiate the answer to obtain the solution

$$S(t) = S(0) \exp \left\{ \int_0^t \sigma(s) dW(s) + \int_0^t \left( \alpha(s) - \frac{1}{2} \sigma^2(s) \right) ds \right\}.$$

**4.13** Suppose  $B_1(t)$  and  $B_2(t)$  are Brownian motions and

$$dB_1(t) dB_2(t) = \rho(t) dt,$$

where  $\rho(t)$  is a stochastic process taking values strictly between  $-1$  and  $1$ . Define processes  $W_1(t)$  and  $W_2(t)$  such that

$$\begin{aligned} B_1(t) &= W_1(t), \\ B_2(t) &= \int_0^t \rho(s) dW_1(s) + \int_0^t \sqrt{1 - \rho^2(s)} dW_2(s), \end{aligned}$$

and show that  $W_1(t)$  and  $W_2(t)$  are independent Brownian motions.